

فحوصات المواد المخدرة

Drug Testing



biolab
مختبر بيولاب الطبي



فحوصات المواد المخدرة

الأدوية المشروعة: هي الأدوية التي يسمح بصرفها من قبل الطبيب وتكون متوفرة ومتاحة في الصيدليات.

الأدوية الغير مشروعة: هي الأدوية الغير مسموحة قانونياً بالبيع والتداول وتسبب مخاطر كثيرة منها تلف الدماغ، القلب وغيرها.



لماذا يلجأ البعض لاستعمال المخدرات غير المشروعة؟

قد يلجأ الشخص للمخدرات غير المشروعة لأسباب عديدة:

• للهروب من الواقع لفترة من الوقت

• الاكتئاب

• قد يساعد الشخص مؤقتاً بالشعور على نحو أفضل أو نسيان المشاكل

• البحث عن الشعور الذي قد يحصل للشخص من بعد استعمال المخدرات

الشباب (المراهقين) قد يستعملون المخدرات للأسباب التالية:

• للشعور بالنضج

• المجارة الاجتماعية

• للاسترخاء والشعور بالرضا

• لاتخاذ المخاطر و التمرد

• الارضاء الفضول



هل يمكن الاستدلال على متعاطي المخدرات؟

متعاطي المخدرات قد تظهر عليهم
بعض الأعراض المذكورة أدناه:

- تقلب المزاج، السلبية والقلق طوال الوقت
- يفضل الوحدة كثيراً
- صعوبة في التركيز
- النوم الكثير
- حب المشاكل والقتال
- حمرة العينين وتنفخهم
- الخسارة أو زيادة الوزن
- السعال
- سيلان الأنف طوال الوقت



ما هي الفحوصات التي يمكنني إجرائها؟

فحص المخدرات في **البول** هو الأكثر شيوعاً في الكشف عن تعاطي المخدرات، ويمكن اكتشاف المخدرات حتى لو تم التعاطي في الاونة الأخيرة أو حتى لمرة واحدة.

وقت الكشف عن المخدرات هو الفترة الزمنية التي يمكن الكشف فيها عن المخدرات في الجسم، و يتأثر هذا الوقت بتكرار عدد مرات التعاطي وطريقة الاختبار وعمليات أيض الجسم وغير ذلك من العوامل.

تفسير نتائج الفحص

سلبى	النتيجة الطبيعية
إذا كانت نتيجة الاختبار ايجابية لواحد من المخدرات فإن ذلك يعني أن الكشف عن المخدرات في البول ايجابي.	النتيجة غير الطبيعية

Type of Drug	Proprietary or street name	Medical uses	Rout of Administration	Durg Detection time
Cocaine	Coke, Crack, Flake, Rocks, Snow	Local anesthetic vasoconstrictor	Injected, smoked, sniffed	1- 4 days



Cannabinoids (Marijuana, Hashish)	Blunt, Grass, Herb, Pot, Reefer, Sinsemilla, Smoke, Weed	None	Oral, smoked	1 day - 5 weeks
Amphetamine	Biphphetamine, Dexedrine; Black beauties, Crosses, Hearts	Attention deficit hyperactivity disorder (ADHD), obesity, narcolepsy	Injected, oral, smoked sniffed	1- 2 days
Barbiturates	Amytal, Nembutal, Seconal, Phenobarbital ; Barbs	Anesthetic, anticonvulsant, hypnotic, sedative	Injected, Oral	2 - 10 days
Benzodiazepines	Ativan, Halcion, Librium, Rohypnol, Valium; Roofies, Tranks, Xanax	Antianxiety, anticonvulsant, hypnotic, sedative	Injected, Oral	1- 6 weeks



Methadone	Amidone, Dolophine, Methadose	Analgesic, treatment for opiate dependence	Injected, Oral	1 day- 1 week
Methamphet amine	Desoxyn; Crank, Crystal, Glass, Ice, Speed	ADHD, obesity, narcolepsy	Injected, Oral, smoked, sniffed	1- 2 days
Morphine	Roxanol, Duramorph	Analgesic	Injected, Oral, smoked	1- 2 days
Phencyclidine	Angel dust, Supergrass, Boat, TicTac, Zoom, Shermans	Anesthetic (veterinary)	Injected, Oral, smoked	2 - 8 days
Tricyclic Antidepressant	Amtriptyline(Trypti zol, Domicol, Lentizol, Limbitrol, Triptafen) Colmipramine (Anafranil) Imipramine (Tofranil) Lofepramine (Gamanil)	Major depressionand in lower doses for insomnia and pain relief in some chronic pain syndromes.	Injected, Oral	Depends on the type of antidepressant



ماذا يمكن أن أفعل؟

التخلي عن ادمان المخدرات أمر صعب ولكنه ليس مستحيلاً. إذا كنت تعتقد أن أحدهم يتعاطى المخدرات، ان أفضل شيء يمكن أن تفعله هو أن تخبر أحد الباغين ذو ثقة. قد يكون أحد الوالدين أو غيرهما من الأقرباء أو المدرس أو مدرب الرياضة أو مستشار المدرسة. قد يحتاج الشخص المتعاطي مساعدة خاصة للتوقف عن تعاطي المخدرات، العائلة والأصدقاء يمكنهم أن يلعبوا دوراً حاسماً وهاماً في تحفيز متعاطي المخدرات بدء العلاج وإكمالها ويمكنهم تعزيز وتوسيع نطاق فوائد العلاج.







Drug Testing

What can i do about it?

Breaking a drug addiction is difficult, but not impossible. If you think someone is using drugs, the best thing to do is to tell an adult that you trust. This could be a parent, other relative, teacher, coach or school counselor.

The person might need professional help to stop using drugs.

Family and friends can play critical roles in motivating individuals with drug problems to enter and stay in treatment and can strengthen and extend treatment benefits.



Methadone	Amidone, Dolophine, Methadose	Analgesic, treatment for opiate dependence	Injected, Oral	1 day- 1 week
Methamphet amine	Desoxyn; Crank, Crystal, Glass, Ice, Speed	ADHD, obesity, narcolepsy	Injected, Oral, smoked, sniffed	1- 2 days
Morphine	Roxanol, Duramorph	Analgesic	Injected, Oral, smoked	1- 2 days
Phencyclidine	Angel dust, Supergrass, Boat, TicTac, Zoom, Shermans	Anesthetic (veterinary)	Injected, Oral, smoked	2 - 8 days
Tricyclic Antidepressant	Amtriptyline (Trypti zol, Domicol, Lentizol, Limbitrol, Triptafen) Colmipramine (Anafranil) Imipramine (Tofranil) Lofepramine (Gamanil)	Major depression and in lower doses for insomnia and pain relief in some chronic pain syndromes.	Injected, Oral	Depends on the type of antidepressant



Cannabinoids (Marijuana, Hashish)	Blunt, Grass, Herb, Pot, Reefer, Sinsemilla, Smoke, Weed	None	Oral, smoked	1 day - 5 weeks
Amphetamine	Biphphetamine, Dexedrine; Black beauties, Crosses, Hearts	Attention deficit hyperactivity disorder (ADHD), obesity, narcolepsy	Injected, oral, smoked sniffed	1- 2 days
Barbiturates	Amytal, Nembutal, Seconal, Phenobarbital ; Barbs	Anesthetic, anticonvulsant, hypnotic, sedative	Injected, Oral	2 - 10 days
Benzodiazepines	Ativan, Halcion, Librium, Rohypnol, Valium; Roofies, Tranks, Xanax	Antianxiety, anticonvulsant, hypnotic, sedative	Injected, Oral	1- 6 weeks



How can i test for it?

Drug screening in **urine** is the most common test used for detecting drug abuse because it's less invasive than blood analysis, typically the least expensive and can detect infrequent or recent single use.

Drug Detection Time is the time period in which the drug tests positive in your system. It is affected by frequency of use, type of drug, test method, metabolism and other factors.

Result Interpretation

Normal result	Negative
Abnormal result	If the test result is positive for one of the drugs, it means that the drug is detected in urine. In some cases, a test will register a false positive result, this can occur from interfering factors such as, some food or prescription medications.

Type of Drug	Proprietary or street name	Medical uses	Rout of Administration	Durg Detection time
Cocaine	Coke, Crack, Flake, Rocks, Snow	Local anesthetic vasoconstrictor	Injected, smoked, sniffed	1- 4 days



How can i tell if someone is using drugs?

A person using drugs may have some of the symptoms mentioned below:

- Become moody, negative or worried all the time
- Ask to be left alone a lot
- Have trouble concentrating
- Sleep a lot
- Get in fights
- Have red or puffy eyes
- Lose or gain weight
- Cough a lot
- Have a runny nose all of the time



Why do people use illegal drugs?

A person may use illegal drugs for many reasons:

- To escape from reality for a while
- Depression
- It may temporarily help a person feel better or forget about problems
- People seek the feeling they might get after using drugs

Young people (teenagers) may use drugs for the following reasons:

- To feel grown up
- To fit in belong
- To relax and feel good
- To take risks and rebel
- To satisfy curiosity



Drug Testing

Drugs are chemicals that change the way a person's body work; there are two types of drugs:

1- Legal drugs : Medicines that are allowed to prescribe from doctors to patients and are available in the pharmacies.

2- Illegal drugs : Are drugs which have limitations on their ownership and they can damage the brain, heart and other important organs.



فحوصات المواد المخدرة

Drug Testing



biolab
مختبر بيولاب الطبي

